

| CUSTOMER INFORMATION | | SAMPLE INFORMATION | | |
|----------------------|----------------------|--------------------|----------|--|
| Name: | Josh Toal | Grower Name: | RhizeBio | |
| Email | support@rhizebio.com | Field ID: | Field 1 | |
| Phone | 919-351-2318 | Product Type: | Compost | |
| Bill To: | RhizeBio | Feedstock(s): | N/A | |

| Test ID/Product Name | RhizeBio Compost #1 | Test Date | 3/26/2025 |
|----------------------|---------------------|-----------|-----------|
|----------------------|---------------------|-----------|-----------|

| Species Summary | Functions of Interest | | | Top 20 Genera Identified | | | |
|---------------------------------------|-----------------------|-------|--|--------------------------|------------------|--|----------|
| Microbial Community Composition | | | Oxygen Availability | | | AE = Aerobic AN = Anaerobic | |
| Number of Microbes and Viruses 1,482 | | 82 | Anoxic Environment 44% Medium | | Medium | OA = Obligate Aerobic ON = Olbigate Anaerobic F = Facultative MA | |
| Top 10 Percentage | 48% | | High Oxygen Environment | 44% | Medium | = Microaerophilic | |
| Top 50 Percentage | 73% | | Low Oxygen Environment | 2% | Very Low | 1 Streptomyces | 8.37% F |
| Top 100 Percentage | 81% | | Carbon Cycling | | | 2 Bradyrhizobium | 6.23% MA |
| Bacteria to Fungal Ratio | 297 to 1 | | Organic Carbon Breakdown | 76% | High | 3 Nocardioides | 5.18% AN |
| Ectomycorrhizal to Arbuscular Ratio | | to 1 | Methanogenesis | 83% | High | 4 Pseudomonas | 4.53% AN |
| Viruses and Pathogens | | | Carbon Fixation | 55% | Medium | 5 Burkholderia | 3.00% AN |
| Total Virusesand Percentage | 11 | 0.75% | Nitrogen Cyclin | Nitrogen Cycling | | 6 Mycolicibacterium | 1.81% |
| Pathogens Percentage - Top 100 | 0 | 0.00% | Denitrification | 2% | Very Low | 7 Arthrobacter | 1.61% AN |
| Microbial Metabolism | | | Nitrification | 75% | High | 8 Microbacterium | 1.54% AN |
| Total Bacterial Genera and Percentage | 1434 | 96.8% | Nitrogen Fixation | 88% | Very High | 9 Variovorax | 1.52% AN |
| Anaerobic Bacteria Genera | 245 | 16.5% | Organic Nitrogen Breakdown | 61% | Medium | 10 Massilia | 1.46% AN |
| Aerobic Bacterial Genera | 801 | 54.0% | Nutrient Cycling | | 11 Sphingomonas | 1.44% AN | |
| Unclassified Bacterial Genera | 388 | 26.2% | Phosphorus Mobilization | 37% | Low | 12 Mesorhizobium | 1.43% |
| Total Fungal Genera and Percentage | 37 | 2.50% | Potassium Solubilization | 60% | Medium | 13 Micromonospora | 1.34% AN |
| Anaerobic Fungal Genera | 4 | 0.27% | Sulfur Oxidation | 8% | Very Low | 14 Mycobacterium | 1.33% AN |
| Aerobic Fungal Genera | 28 | 1.89% | Sulfur Reduction | 50% | Medium | 15 Rhizobium | 1.14% AE |
| Unclassified Fungal Genera | 5 | 0.34% | Calcium Transport | 67% | High | 16 Paenibacillus | 1.04% AE |
| Key Metabolic Functions | | | Iron Acquisition | 56% | Medium | 17 Rhodococcus | 0.89% AN |
| Facultative Anaerobic Organisms | 95 | 6.4% | Other Plant Growth Promoting Functions | | 18 Amycolatopsis | 0.86% | |
| Obligate Anaerobic Organisms | 17 | 1.1% | Nodulating Bacteria | 20% | Very Low | 19 Telluria | 0.81% AE |
| Microaerophilic Organisms | 45 | 3.0% | Plant Stress Adaptation | 34% | Low | 20 Conexibacter | 0.80% AE |

RhizeBio, Inc.[™] Test Protocol V1 May 17, 2024